Memoirs of the Queensland Museum.

VOL. X, PART III.
Issued 14th March, 1933.

EDITED BY THE DIRECTOR, HEBER A. LONGMAN, F.L.S., C.M.Z.S.

ISSUED BY THE AUTHORITY OF THE CHIEF SECRETARY FOR QUEENSLAND, THE HON. W. FORGAN SMITH.

CONTENTS.

PAGE.

A New Dinosaur from the Queensland Cretaceous—Plates XV-XVII, Text-figures 1–3. – – – – – Heber A. Longman – – – 131–144

Notes on Australasian and Melanesian Dermaptera. Text-figures 1–4. – – – – – – – – Morgan Hebard – – – 145–156

A NEW DINOSAUR.

From the Queensland Cretaceous.

BY HEBER A LONGMAN, DIRECTOR, QUEENSLAND MUSEUM.

(Plates XV–XVII, Text-figures 1–3.)

INTRODUCTION.—When visiting "Clutha" Station, Maxwelton, North-west Queensland, in August last, Dr. M. J. McKillop, of Brisbane, was interested in some large fossil remains of vertebræ which were shown to him by his brother, Mr. H. J. McKillop, the manager. In response to my request, through Dr. McKillop, these specimens were carefully packed and forwarded to the Queensland Museum, where they were received on 9th January, 1933.

LOCALITY.—"Clutha" is north of the Flinders River, and the homestead is 34 miles from Maxwelton, on the Great Northern Railway, 337 miles west of Townsville, North Queensland.



Text-figure 1.—Part of Clutha Station showing Whitewood Paddock. Site of Fossils marked X.

Del. T. C. Marshall, from Qld. Four Mile Maps, Sheet 82 (Enlarged).

Mr. H. J. McKillop informs me that the fossils were first discovered by the overseer, Mr. H. B. Wade, who noted them on the surface partly exposed in black soil on open down country, "in what is known as Whitewood Paddock, at a spot approximately three miles west of the homestead and about

two miles from Coolibah Creek. There are no public roads on 'Clutha,' but there are many private tracks on the run, and one of these passes within a few hundred yards of the site of the discovery. The area of the paddock is 1,200 acres." (Text-figure 1.)

As the locality is within a district that has provided many fossils representing Lower Cretaceous marine reptiles, it was anticipated that the large vertebræ would be those of *Kronosaurus queenslandicus*,[1] the largest known species. To my surprise and intense interest, however, when the "Clutha" fossils were examined it was obvious that they represented a new giant Dinosaur, the first to be obtained from the Lower Cretaceous deposits of Queensland. The stratigraphical problems associated with this discovery are discussed later.

AUSTROSAURUS* McKILLOPI, genus and species new.

Dorsal vertebræ markedly opisthocœlous; centra with thin cortical walls, much enlarged at the enarthrodial articulations; intramural region a complex of small cavities; pleurocœles prominent, with external and internal divisions. Neural arch with deep recess between the prezygapophysial lamina and the infradiapophysial buttress.

This tentative generic diagnosis will obviously need to be supplemented when more complete material is forthcoming. The proportions and some of the characteristics of the vertebræ, however, appear significant, and it is anticipated that this Australian Cretaceous Dinosaur will prove, when better known, as distinctive as most of the giant Sauropoda found in similar deposits elsewhere.

Material.—This consists of three massive blocks, each containing incomplete paired vertebræ cemented together at the enarthrodial articulations of the centra. Owing to the extreme development of the intramural cavities, these vertebræ, although so massive, are very fragile. With the exception of the articulating surface of the "cups" and "balls," the peripheral walls are surprisingly thin. The layers of bone have now been strengthened by repeated soakings in shellac solution. The matrix (described elsewhere) is much more durable than the actual fossil, and the partial clearing of the bones necessitated much patient work. The specimens were evidently weathered out of the original formation long before their discovery, and they were found lying in

[1] Longman, H. A., Mem. Qld. Mus., VIII, Pt. 1, 1924, X pt. 1, 1930, and X pt. 2, 1932.

* Following numerous precedents, the prefix "Austro" has been used for its present geographical significance, and this is also in consonance with the nomenclature of many modern genera of Dinosaurs, distinctively named from other parts of the world.

black soil, some rootlets actually being present in the interstices. Owing to the effects of prolonged abrasion and weathering, large portions of the original contours have been lost.

For convenience in description the specimens have been designated A, B, and C. (Reg. No. F2316.)

Description.—Specimen A consists of the major portion of one centrum, with the lower part of the neural arch, and the fractured "cup" of the contiguous unit, which has fragments of its arch in the matrix above. In the more complete vertebra, the posterior part of the centrum has been shorn away, and the cross section of fracture shows within the thin walls a coarse mosaic of irregular cancelli enclosed by thin bony plates. These intersecting laminæ form a labyrinth throughout the exposed transverse section of the centrum, but are most prominent where supporting the cortical walls. The cancelli are wholly or partly infilled with a crystalline matrix of carbonate of lime (described elsewhere).

The lateral walls of the right side of the centrum have disappeared, but the anterior part of the left lateral surface is preserved, with an almost complete pleurocœle. This area has been largely freed from the matrix and exhibits important characteristics (Plate XV.). Of the actual walls of the inferior surface of the centrum only a small portion is preserved, but the general proportions are apparent. When viewed from below, it is seen that the body of the centrum is constricted and the lateral walls curve outwards anteriorly to the much larger rim of the "ball." This constricted portion of the centrum, with its associated lateral cavities, is evidently fragile in comparison with the articular areas, for all of the specimens are fractured in this region.

The dimensions of Specimen A are as follows:—Maximum length $10\frac{1}{2}$ inches (267 mm.); height $16\frac{1}{4}$ inches (413 mm.); transverse diameter $10\frac{1}{8}$ inches (257 mm.).

Pleurocœle.—This is situated in the upper portion of the centrum and is well shown in Plate XV. This large lateral cavity is divided into an external and an internal portion. The former is a large horizontal oval recess, from which the matrix has been cleared, and at the bottom is a much smaller oval fenestra, the internal pleurocœle, which extends further into the centrum. The external recess is about 42 mm. in depth; its antero-posterior diameter is approximately 90 mm., and its vertical height is 60 mm. The lower wall of this cavity forms an almost horizontal shelf, but, from the inner surface, the upper and anterior walls curve gradually outwards to the plane of the lateral surface. The matrix has been almost entirely cleared from the internal pleurocœle, the diameters of which are approximately 50 mm. by 30 mm. This inner recess is divided into two portions by a thin vertically-placed bar of bone, situated at the anterior third.

On the right side the walls of the centrum have entirely disappeared. The pleurocœle is shown in section, and was originally infilled with matrix which was surrounded with very thin peripheral walls. The typically marine matrix has now been removed on each side, so far as it is safe to do so. The two pleurocœles are not confluent, but the matrix terminates against the laminar complex, which forms so distinctive a feature of the centrum, within about 20 mm. from the median line. No trace of a median vertical partition can be seen in the fractured surface across the centrum, and the laminar complex evidently takes the place of this structure.

Neural Arch.—This region is very incomplete, and the right side has largely disappeared. The full height is not preserved, but the fractured surface above exhibits a complex of abraded and distorted elements. It is obvious, however, that the laminæ and brackets were very extensive and were lightly framed.

The preserved portions present significant similarities in this region with those shown in the well-known figure of the vertebra of "*Ornithopsis*" *hulkei*, first described and figured by J. W. Hulke in 1880.[1]

Owing to the presence of an almost complete fracture in the fossil, it has been possible to remove the major portion of the neural arch from its basal part. The exposed contours give some interesting details of the internal structure. From the upper margin of the centrum, near the anterior rim, a curved plate, somewhat "saddle-shaped," is produced anteriorly and outwardly. The basal portion of its anterior surface, when viewed from the lateral aspect, arises almost vertically, but the upper portion, as may be seen from impressions in the matrix, expands outwardly and anteriorly, evidently supporting prezygapophyses which projected over the border of the contiguous vertebra (Plate XVI.). This infraprezygapophysial lamina (to use Osborn and Mook's revised term[2]), is very incomplete, but it appears to have been much more extensive in this vertebra than in other related Dinosaurs, judging from impressions in the matrix. Its median portion is incomplete. The base of a similar lamina is preserved on the right side, and this meets the base of the left lamina above the neural canal. When viewed from above, the fractured section shows that the median extension of each lamina curves posteriorly to meet its fellow, forming a wide recess above the neural canal. The hypantrum-hyposphene articulation is not preserved, but was evidently present above the fractured section.

A large recessed area lies between this anterior lamina and the infra-diapophysial buttress. This recess is relatively very large and attains a depth of at least 60 mm. (Text-figure 3). Only the basal part of the median portion

[1] Hulke, J. W., Quart. Journ. Geol. Soc., Vol. 36, pp. 32-34, plate iv.

[2] Osborn and Mook, Mem. Amer. Mus. Nat. Hist, n. s. III., 1921.

Austrosaurus mckillopi. Lateral view of Specimen A—incomplete dorsal vertebræ (Holotype).

(*Approximately one-third natural size.*)

Face page 134.

Photo., W. J. Sanderson.

of the pedicle of the arch is preserved, as may be seen in Plate XV. Unfortunately the postzygapophysial region and the whole of the platform supporting the transverse process are missing. Judging from fragments exposed in the matrix over the remnant of the contiguous vertebra, however, the region of the diapophyses was situated high in the neural arch,—at least as high above the centrum as the vertical diameter of the centrum.

Rib.—A fragment of a rib is preserved in the matrix between the two vertebræ on the left side (Plate XV.). In section this is oval, with one side flattened, the diameters being 90 mm. by 40 mm.

Neural Canal.—The matrix in the site of the neural canal is well preserved in all three specimens, and is transversely oval in section. The average transverse diameter is 50 mm., and the vertical height is 35 mm. The cylinders of matrix are partly exposed above in Specimens B (Plate XVII.) and C.

Immediately above the region of the canal in A is an arch of bone showing an elongated isosceles triangle in section, as seen in the matrix, but the plates are very thin as preserved. This structure is seen in the anterior and posterior sections now exposed in Specimen A, and it is also noticeable in the section of Specimen B. It appears to be a distinctive feature of the intramural laminæ of the neural arch (Plates XVI. and XVII.).

Specimen B shows subequal portions of two incomplete vertebræ. The anterior element mainly consists of the expanded rim of the "cup," which is closely adpressed to the "ball" of the adjoining unit. The preserved portion of the body of the centrum is almost entirely intramural and consists of a mosaic of large cancelli. The matrix of part of the neural canal is present.

The associated vertebral fragment consists of an incomplete centrum containing a pleurocœle on each side. On its posterior surface it exhibits the inner portion of the deep cup. So much of the rim is missing, however, that the maximum length of the complete vertebra cannot be precisely estimated. The concavity is now covered with a thin tessellated layer, which evidently represents the original cartilage now changed to crystals of carbonate of lime (Plate XVII.). The preserved portion is 180 mm. in transverse diameter, with a vertical height of 135 mm.

The anterior base only of the neural arch is preserved. As exposed in the matrix, the right and left anterior laminæ exhibit "saddle-shaped" contours resembling those described from the more complete elements in specimen A. The neural canal matrix, consisting of a vertically compressed oval cylinder (54 mm. x 40 mm. in section) is also preserved. Above this an elongated isosceles triangle of matrix is enclosed within thin plates of bone, as in Specimen A.

The pleurocœles in this centrum are less elongated than those in Specimen A, and are infundibulate. They are more obliquely placed in the

upper part of the centrum. The great differences in the size of the pleurocœles shows that B was not consecutive with A in the vertebral series, but it is considered that they were not widely separated.

On the lower surface the peripheral walls are incomplete. In section the contours are broadly convex, but medially there is a shallow depression running longitudinally along the centrum.

The dimensions of Specimen B are :—Maximum length $11\frac{3}{4}$ inches (299 mm.) ; transverse diameter $12\frac{1}{4}$ inches (306 mm.) ; height $12\frac{1}{8}$ inches (303 mm.).

Judging from the proportions of the parts preserved, this vertebra was distinctly shorter than those represented in Specimen A, but was much more massive transversely.

Specimen C consists of the massive rim of a "cup" shorn from a centrum and cemented with matrix to the associated "ball." On the lower surface, portions of the external walls are preserved, although greatly crushed, but otherwise the cortical bone has entirely disappeared. On the right side an oval mass of matrix (60 mm. x 25 mm.) is seen obliquely situated deep in the body of the matrix, denoting the inner part of a pleurocœle. Near the border of the "cup," midway between the inferior and superior margins of the centrum, an additional oval mass of matrix is seen on one side. This matrix, which is a small oval vertically placed, apparently represents an additional pleurocœle, much smaller (diameters approximately 40 x 20 mm.) and quite separate from the larger lateral cavity.

Specimen C has been subjected to great vertical pressure, and the cylinder of matrix representing the neural canal has been crushed down into the cancellous tissue of the centrum. The expanded rim, although cracked, has not suffered so much as the median portions, but the transverse diameter is probably somewhat enlarged by intense pressure during fossilisation.

Specimen C is relatively massive in comparison with the other fragments. Its transverse diameter is no less than $13\frac{1}{4}$ inches (336 mm.), The maximum length is $11\frac{3}{4}$ inches (299 mm.—and the height of the specimen as preserved is 9 inches (229 mm.).

The predominance of the transverse diameter over the vertical diameter—quite apart from the distortion owing to pressure—suggests that the two vertebræ represented came from near the sacral region.

Although very incomplete, these specimens definitely show that this Queensland Cretaceous Dinosaur exhibits in its dorsal series of vertebræ the variability which is characteristic of most of the Sauropoda.

Austrosaurus mckillopi. Posterior view of Specimen A, showing transverse fracture of centrum and intramural complex of cavities.

(*Approximately one-third natural size.*)

Face page 136.

Photo., W. J. Sanderson.

Affinities.—Matley (1931, p. 281),[1] in commenting on the connexion in Mesozoic times of South America with south-eastern Asia by way of Antarctis, the Australian region and the Sunda Archipelago, as discussed by von Huene, says that it is "really remarkable that all the three Cretaceous sauropod genera of India—*Titanosaurus*, *Antarctosaurus*, and *Laplatasaurus*—should be found so far away as Patagonia." &c. Notwithstanding the wide range of these and other sauropodous genera, it is difficult to place this Queensland Cretaceous Dinosaur with any well-known types, in so far as comparisons can be made with our material. *Austrosaurus* does not closely resemble any of the species described by von Huene[2] in his comprehensive work "Los Saurisquios y Ornitisquios del Cretaceo Argentino," and it evidently does not belong to the Family Titanosauridæ.

Austrosaurus mckillopi appears to be a more specialized Dinosaur than the Queensland Jurassic Sauropod *Rhætosaurus brownei*, described by the writer.[3] In *Austrosaurus* the centra of the dorsal vertebræ attain a maximum of specialization in the intramural complex of laminæ enclosing irregular cavities, supplementing the actual recesses of the pleurocœles. It is not clear whether the pleurocœles are actually connected with the complex of small internal cavities. Judging from the condition of the matrix, however, they were not, and probably only the lateral recesses were actually pneumatic in life. Owen's view[4] that the inner recesses of such centra were filled with "chondrine," although opposed by Seeley and Hulke,[5] may well be true.

Apparently there are two types of cavernous dorsal vertebræ in the Sauropodous Dinosaurs. In one type, the lateral cavities or pleurocœles extend into the intramural area and form an enlarged chamber, recessed behind the actual opening: there is no complex of small cavities and the additional required strength of the centrum is formed by thick portions of the peripheral walls and a median vertical partition.

This type of vertebra is illustrated by the transverse section of the trunk centrum of "*Ornithopsis*" figured by Hulke in 1879,[6] here reproduced (Text-figure 2). The section of a dorsal vertebra of Marsh's *Morosaurus lentus* (*Camarasaurus*, Cope) illustrates another development of this type,[7] and many other examples are known. Osborn and Mook record (*loc. cit.*, 1921, p. 306) that in *Camarasaurus* the pleurocœles "occupy most of the bodies of the centra."

---

[1] Matley, C. A., Geol. Mag., Vol. LXVIII, 1931.

[2] Huene, F. von, Anales del Museo de La Plata, tomo III, serie 2, 1929.

[3] Longman, H. A., Mem. Qld. Mus., VIII, pt. 3. pp. 183-194, Plates XXIX-XXXIII; Mem. Qld. Mus., IX, pp. 1-18, Plates I-V; Australian Museum Mag., Vol. III, No. 3, pp. 97-102; Mem. Qld. Mus., IX, pt. 3, p. 249, Plate XXIX.

[4] Owen, R., Mon. Foss. Rept. Weald. & Purb., Supp. No. VI, 1874, p. 6.

[5] Owen, R., Quart. Journ. Geol. Soc., vol. 35, 1879, p. 762.

[6] Hulke, J. W., Quart. Journ. Geol. Soc., 1879, Vol. 35, p. 756.

[7] Marsh, O. C., The Dinosaurs of North America, 16th Ann. Rep. U. S. Geol. Sur., 1896. Plate XXXII, fig. 2a.

In the other type, the lateral cavity or pleurocœle forms a recess which is not expanded within the centrum; the peripheral walls may be surprisingly thin, and most of the intramural space is filled with a complex of small laminar cavities. These laminæ reinforce the thin peripheral walls, exhibiting an extraordinary combination of strength with relative lightness. The majority



Text-figure 2.—Transverse section of dorsal vertebra of *Ornithopsis* (after J. W. Hulke). *n.* Neural canal, *p.* partition, *b.* buttress, *c.* chamber, *o. o.* lateral openings to chambers.

of the small inner cavities appear to be more or less insulated, and there is no evidence of connecting passages between the lateral recess and this inner complex. This type of vertebræ is well illustrated in *Diplodocus carnegii*, as fully described by J. B. Hatcher (1901), who refers to the "remarkable degree of specialization, unsurpassed if not unequalled by other vertebrates in the general characters of its vertebral column in its adaptation of these mechanical principles which combine maximum strength with minimum weight (p. 12).[1]

It is very evident that the vertebræ of *Austrosaurus mckillopi* resemble those of *Diplodocus* in the special structure of the centrum. A transverse section of the centrum of Specimen A, shown in Text-figure 3, illustrates the proportions of the pleurocœles, taken at their maximum extent, and the intramural complex of cavities surrounding them. The recess behind the anterior lamina, above the centrum, is also indicated.

A longitudinal section of a vertebra of *Austrosaurus* would exhibit a somewhat similar structure, but more complex, to that figured by Owen in his "*Chondrosteosaurus*" from the Wealden deposits of the Isle of Wight.[2]

---

[1] Hatcher, J. B., Mem. Carnegie Museum, Vol. 1, 1901, p. 12.

[2] Owen, R., Mon. Foss. Rept. Wealden & Purbeck, Pal. Soc. Supp. VII, 1876, Pl. V, fig. 2.

The type centrum of "*Bothriospondylus magnus*" as figured by Owen, which is also the type of Seeley's *Ornithopsis*, shows the same characteristics.[1] Seeley's and Owen's genera are now usually included in the synonymy of Mantell's

Text-figure 3.—Section of vertebra of *Austrosaurus mckillopi* taken at the maximum extent of the pleurocœles, showing also the recess in the neural arch. The complex of intramural cavities is shown somewhat diagrammatically. (One-half natural size.) *Del. M. J. McKillop.*

[1] Owen, R. (*loc. cit.*), part 2, Plate IX, fig. 1.

Pelorosaurus,[1] but if the centra figured by Hulke, previously mentioned, are congeneric, there is an extraordinary range of variation in structure—even for a sauropodous Dinosaur—in this genus.

A. Smith Woodward, records: "In *Ornithopsis* the centrum of each vertebra is chambered throughout, and the thin partitions between the small cavities consist of hard dense bone.[2]

In his paper, "Zur Systematik und Biologie der Sauropoden" (1930)[3] Baron Franz Nopesa touches on the differential development of pleurocœles, but he does not deal with the significance of an intramural complex of small cavities, and I can find no special references in Baron Huene's papers.

In Zittel's Text-book of Palæontology (Eng. Ed. 1902) only two families of Sauropoda, the Camarasauridæ and the Diplodocidæ, are recognised, whereas Marsh had previously recorded six. Four families are listed in "The Osteology of the Reptiles" by S. W. Williston (edited by W. K. Gregory, 1925), with a long list of genera *incertæ sedis*. Huene, however, lists six families in 1927 ("Short Review of the Present Knowledge of the Sauropoda," These Memoirs, Vol. IX. pt. 1, 1927). He divides the Cetiosauridæ into two sub-families: "Cardiodontidæ and Brachiosauridæ," and points out that the vertebræ of the latter are more cavernous than those of the former, although neither are so cavernous as in the families Morosauridæ and Diplodocidæ. It is doubtful whether the multiplication of families will tend to elucidate the phylogeny of this group. At present we have specialized genera such as *Diplodocus* in the Upper Jurassic with no known descendants in the Cretaceous, whilst, as R. S. Lull and F. Nopesa[4] have pointed out, the comparatively simple type of *Titanosaurus* occurs in the Upper Cretaceous.

As von Huene states, the natural classification of the Sauropoda is no yet clear. Doubtless our knowledge of their phylogeny will increase with new and more complete material. Where specimens are fragmentary the allocation of certain genera to families depends on the significance attached to certain features. As R. S. Lull points out, *Barosaurus* has several features in common with *Diplodocus* (including the two-branched characteristics of the chevrons, once thought distinctive), yet these are placed in distinct families.[5]

Notwithstanding the striking similarity between the intramural complex of the centra in *Austrosaurus* and *Diplodocus*, this Queensland Cretaceous

---

[1] Mantell, G. A., Phil. Trans. Roy. Soc., 1850, p. 379.

[2] Woodward, A. Smith, P.Z.S., 1905, p. 232.

[3] Nopcsa, F., Palæobiologica, iii., Band, 1930.

[4] Nopcsa, F., Quart. Journ. Geol. Soc., Vol. LXXIX, 1923, p. 107.

[5] Lull, R. S., Mem. Connect. Acad. Vol. VI., 1919, p. 40.



Austrosaurus mckillopi. Posterior view of Specimen B.
(*Four-ninths natural size.*)

Face page 140.

Photo., W. J. Sanderson.

Dinosaur is tentatively placed in the family Cetiosauridæ, as a specialized member, allied to *Pelorosaurus*. As Riggs' Brachiosauridæ was primarily established on the relative dominance of the fore limb, additional evidence will be required before it can be affined with this series.[1]

Austrosaurus does not appear to be closely related to either *Cetiosaurus*, *Cetiosauriscus*, or *Haplocanthosaurus*, which are placed by von Huene in the sub-family Cardiodontidæ. Hatcher[2] regards the last-named genus as representing "the most primitive Sauropod known" (p. 45), but he records that the "vertebræ show numerous large intramural cavities instead of the close, though cancellous texture" of *Cetiosaurus* (p. 53).

With the few fragments available, the author is diffident in doing more than suggest, at present, the affinities of this new Cretaceous Dinosaur. Attempts to solve problems of phylogeny can be best made by those who have had opportunities of examining the mass of material stored in European and American Museums. It is believed, however, that the intramural complex of the vertebral centra exhibits diagnostic characters to which greater attention should be given. The greatest development of what may be called the cryptocamarillan type is found in the posterior dorsals, whilst the phanerocamerate or open-chambered type of centra obviously reaches its maximum of evolution in the cervicals of specialized species.

Many genera that are now well known were first described from very fragmentary specimens. In view of the special interest attached to this great group of Sauropodous Dinosaurs and the study of their rapid evolution throughout Jurassic and Cretaceous times, it is hoped and anticipated that supplementary material will be forthcoming of this giant Queensland reptile.

Additional interest is given to the study of these giant reptiles by the point raised by H. Fairfield Osborn, who considers that the Upper Jurassic Lower Cretaceous migrations of the Sauropoda, probably from a Central Asiatic stock, into all the continents, shed light on the adaptive radiation of mammals.[3]

Dimensions.—Taking the anterior part of the sacrum as representing the central part of the vertebral series, as is characteristic of most of these giant Sauropodous Dinosaurs, and taking the average of dorsal and cervical vertebræ for the group as being twenty-five, a tentative estimate of the length of one of these great reptiles can be made from a single vertebra. In a Dinosaur of the *Brachiosaurus* type excavated by F. W. H. Migeod for the British Museum at Tendaguru in 1930, the length of the dorsal vertebræ varied from nine to fourteen inches, and allowance must be made for regional variation.[4]

---

[1] Riggs, E. S., Field Columbian Museum, Geol. Ser., Vol. II, No. 6, 1904.
[2] Hatcher, J. B., Mem. Carnegie Mus., II., No. 1, 1903.
[3] Osborn, H. Fairfield, Rep. Brit. Assn., 1931, p. 389.
[4] Migeod, F. W. H., Natural History Magazine, Vol. III, No. 19, 1931, p. 92.

Obviously the elongation of the cervicals must be considered, and an unusually long neck, or an attenuated and extended tail would be disturbing factors. It is permissible, however, to make a tentative estimate on the average proportions of well-known forms, and on this basis it is considered that the vertebræ of *Austrosaurus mckillopi* represent a giant Dinosaur of about fifty feet in length, and it probably exceeded in size the large Dinosaur *Rhætosaurus brownei* described by the writer from Jurassic deposits at Durham Downs, South-western Queensland.

Habits.—As the Sauropodous or lizard-footed Dinosaurs are semi-aquatic quadrupeds which were undoubtedly herbivorous in habits, the discovery of these fossils in marine strata, infilled with typically marine matrix and definitely associated with marine shells (*Inoceramus* and *Beudanticeras*), is of considerable interest. Dinosaurs are typically found in fresh-water or estuarine deposits, but there are several exceptions. In the bone beds at Tendaguru, East Africa, the remains of Dinosaurs were found in close association with marine fossils. J. Parkinson (1930)[1] gives interesting details of excavating fossils "showing how close the Dinosaurs lived to the sea at Tendaguru."

It is generally recognised that these huge Sauropods were adapted for wading, with a lengthy, relatively light and flexible neck associated with massive limbs and more solid tail. It is unlikely that they were good swimmers or that they even resembled the modern crocodile in occasionally making lengthy trips in the ocean. But doubtless they roamed around the shores of the shallow Cretaceous sea, as well as frequenting the fresh-water lagoons, swamps, and rivers, feeding on the luxuriant vegetation of the period. It seems probable that the relatively buoyant body of a Dinosaur, after disablement or death, might be carried miles out to sea, providing a feast for such formidable swimming marine reptiles as *Kronosaurus*, until its disassociated bones sank to rest in the limey mud. But whatever the cause of its death and deposition, the presence of these bones in marine beds at "Clutha" is definite evidence of the existence during its lifetime of fresh-water Cretaceous formations nearby, which provided a suitable environment for these giant Dinosaurs.

Matrix.—I am indebted to Dr. F. W. Whitehouse for the following note:—"The matrix consists of a fine-grained, argillaceous limestone (or calcareous mudstone) of a light fawn colour—a rock type common throughout the Cretaceous beds of the Great Artesian Basin. This matrix is found infilling the open cavities of the neural arch, the neural canal, the lateral recesses of the centra, and it also occurs between the vertebræ. Within the inner recesses of the fossil similar matrix is present adjacent to the open cavities. In general, however, these inner cavities are only partly filled and then with coarsely crystalline calcite. The condition thus is similar to that of the preservation of the ammonites in these Cretaceous beds where the general matrix of the beds

[1] Parkinson, John, "The Dinosaur in East Africa," p. 119 (Witherby).

fills in the body chamber; but the earlier and sealed chambers contain coarsely crystalline calcite that rarely fills the whole space. Evidently the calcareous muds filled the large open cavities when the fossil was embedded, but was unable to gain access to the sealed cavities. Percolation of water at a later date through these calcareous beds has apparently been responsible for the deposition and crystallisation of calcite within the inner cavities of the fossils."

THE AGE OF THE BEDS.

Dr. F. W. Whitehouse, Department of Geology, University of Queensland, and Hon. Palæontologist on our staff, kindly contributes the following:—"The matrix, as mentioned above, is typical of the Cretaceous beds of the Great Artesian Basin. These beds consist of a lower or marine suite and an upper or freshwater series. The marine beds, the "Rolling Downs Formation," have been divided into two series:—The Roma Series of Aptian age, succeeded by the Tambo series of Upper Albian age.

"Along the northern railway line from Hughenden almost to Cloncurry all the exposures are in beds of the Tambo Series. North and north-east of these beds, in the eastern portion of the region, beds of the Roma Series appear from below the Tambo Series, forming a marginal fringe to this part of the Cretaceous basin. Beds of this age, yielding Cretaceous ammonites, occur for example at Glendower Station (north-north-east of Hughenden) and Cambridge Downs (north of Richmond). No detailed mapping of the Lower Cretaceous horizons has been done in this area. 'Clutha,' the locality for these fossils, although north of the railway line, from its position might be expected to be in the region of the Tambo Series. This is confirmed by the occurrence of other fossils in the matrix surrounding the bones. This has yielded *Beudanticeras* sp., and fragments of *Inoceramus*. *Beudanticeras* is restricted to the Albian; while in Australia *Inoceramus* is the commonest genus of the Tambo Series, but is almost entirely absent from the Roma Series.

"We may conclude therefore that the beds from which the fossils were obtained belong to the Tambo Series of Upper Albian age."

Mr. L. C. Ball, Chief Geologist, Queensland Geological Survey, has kindly drawn my attention to A. Gibb Maitland's traverses (*see* G.S.Q. Publication No. 121, published in 1898), as bearing on the stratigraphy of the locality. Maitland passed within a few miles to the north of "Clutha" over monotonous open downs entirely devoid of sections.

Previous Records of Dinosaurs.—The only previous record of a Cretaceous Dinosaur in Australia is the report by Sir Arthur Smith Woodward of a "tooth and a posterior caudal vertebra of a small Megalosaurian" from "the Upper Cretaceous opal-bearing sandstone of Lightning Ridge, near Walgett, New South Wales," in 1909.[1] The same author described and figured "an

[1] Woodward, A. Smith, Rep. Brit. Assn., 1909, p. 482.

ungual phalange of a carnivorous Dinosaur," from the Lower Jurassic, Cape Patterson, Victoria, which was unnamed.[1] In 1891 H. G. Seeley described *Agrosaurus Macgillivrayi*,[2] a Saurischian reptile from the north-east coast of Australia, believed to have been collected by Macgillivray from "some locality which was then unnamed," during the voyage of the "Fly." These fossils, consisting of a tibia and a few other fragments indicating "an animal about as large as a sheep," were doubtfully attributed to the Trias. *Rhatosaurus brownei* (Family Camarasauridæ) from the Walloon Series, Jurassic, Durham Downs, South-western Queensland, was first described in these Memoirs by the writer in 1926, with important supplementary descriptions in 1927 and references in 1929.

Acknowledgments.—It is my pleasant duty to thank very heartily Mr. H. J. McKillop for forwarding and presenting this valuable material, Mr. H. B. Wade, who first discovered the fossils, and Dr. M. J. McKillop, of Brisbane, for his kindly interest and also for his skilful sectional drawing of the holotype vertebra.

[1] Woodward, A. Smith, Ann. Mag. Nat. Hist. (7), Vol. 18, 1906, p. 3.

[2] Seeley, H. G., Quart. Journ. Geol. Soc., Vol. 47, 1891, p. 164.

EXPLANATION OF PLATES.

PLATE XV.

Austrosaurus mckillopi. Lateral view of Specimen A—incomplete dorsal vertebræ (Holotype).

PLATE XVI.

Austrosaurus mckillopi. Posterior view of Specimen A, showing transverse fracture of centrum and intramural complex of cavities.

PLATE XVII.

Austrosaurus mckillopi. Posterior view of Specimen B.

NOTES ON AUSTRALASIAN AND MELANESIAN DERMAPTERA.

By Morgan Hebard.

(Text-figures 1–4.)

The following studies are based on the last collections of this Order belonging to the author which have not already been studied. The most important series in it are a small but interesting collection made at Kuranda, Queensland, by F. P. Dodd and a number of specimens received in exchange from the Burr Collection, the data on which has either not been fully published or in some cases not published at all; while some of the determinations are incorrect. To this material has been added a small but very important unstudied collection belonging to the Australian Museum at Sydney, New South Wales. A total of twenty-seven species are treated, of which two are described as new.

In recent years little has appeared on the Dermaptera of the regions here treated, Mjoberg's studies in 1913 and 1924 being among the most important. These are evidently that author's first efforts on this difficult Order where there is still sufficient confusion existing to show plainly why a number of species were incorrectly placed generically by Mjoberg, this and the great individual variation which occurs in many species accounting for a number of synonymic names proposed. Several interesting species were there described, however, and valuable data given on other known forms.

PYGIDICRANIDÆ.

Tagalina semperi Dohrn.

Mt. Lamington, Northern Division, Papua, V, 1927 (C. T. McNamara), 1 large juv. ♀.

In this specimen the pronotum has two dark bands which are convex-convergent to meet meso-caudad and which reach the cephalic margin laterad, that margin being as broadly dark brown. The head is very dark brown except for the pale dorsal and caudal portions of the occiput.

In an adult male from Friederich-Wilhelmshafen, New Guinea, in the author's collection, merest traces of such pronotal marking are present.

Dicrana daemeli (Bormans).

Cairns, Queensland, Australia, (J. F. Illingworth), 1 juv. ♂.

Echinosoma yorkense Dohrn.

Mt. Lamington, Northern Div., Papua, V and VII, 1927, (C. T. McNamara), 2 ♂, 1 ♀. Kuranda, Queensland, 1, 3 and 9, 1925, (F. P. Dodd), 1 ♂, 1 ♀.

There is also a similar male from Waigana, Milne Bay, New Guinea, taken September 9 to 14, 1912 by Carson in the author's collection, received in exchange from the Burr Collection.

We find that *sumatranum* and *yorkense* are variable in coloration and the characters which have been used to separate them can not always be trusted. A minute pale tegminal spot is present in all but the Mt. Lamington female, though very faint in the Kuranda female. Such a spot, believed to be present in *yorkense* but absent in *sumatranum*, is even more distinctly present in a male of *sumatranum* from the Tengger Mountains, Java. Males are, however, easily separated by the abdominal tergites, the fifth to eighth inclusive being laterally heavily keeled and acutely produced in *yorkense*, these roundly produced laterad with only the fifth and sixth weakly keeled in *sumatranum*.

LABIDURIDÆ.

Titanolabis colossea (Dohrn).

Barrington Tops, New South Wales, Australia, I, 13 and 19, 1927, (T. G. Campbell), 5 ♂ (three with total length about 31, two about 39 mm.), [Austral. Mus.] Cutlers Pass, Williams River, N.S.W., X, 28, 1922, (Musgrave and Campbell), 2 small juv., [Austral. Mus.] Hazelbrook, Blue Mountains, N.S.W., IX, 1922, (L. Abrahan), 1 ♀ (total length 34·5 mm.). Blue Mountains, N.S.W., II, 20, 1910, 1 ♀. Booloombayt, Nyall Lakes, N.S.W., IX, 2, 1922, (A. Musgrave), 1 small juv.

A male labelled "N.S.W." is in the author's collection the total length of which measures 53·6 mm.

Gonolabis pacifica (Erichson).

Bairndale, Victoria, (Spry), 1 ♂ (stenolabic), determined by Burr. Barrington Tops, New South Wales, I, 19, 1927, (T. G. Campbell), 1 ♂ (moderately platylabic), [Austral. Mus.) Cutlers Pass, Williams River, N.S.W., X, 28, 1926, (Musgrave and Campbell), 2 ♀, [Austral. Mus.].

We do not consider the characters given for *Mongolabis* Zacher sufficient to warrant recognition of that genus.

Bormans described *brunneri* but stated that it was possibly the same as *pacifica*. Burr in 1908 gave briefly the features which distinguish these species.

Gonolabis tasmanica (Bormans).

Mt. Wellington, Tasmania, 4,000 feet, 1 ♂, 1 ♀, determined by Burr. Interlaken, Tasmania, I, 24, 1928, (A. Musgrave), 1 ♂, [Austral. Mus.].

This insect was placed by Kirby in *Anisolabis* in 1904. The presence of vestigial tegmina afforded Burr's sole reason for referring it to *Euborellia* in 1912. Its agreement with other species of *Gonolabis* is actually very evident.

Anisolabis xenia (Kirby).

Norfolk Island, 1 ♂, 1 ♀.

Nala lividipes (Dufour).

Nepean River, New South Wales, I, 1905, (A. Musgrave), 1 ♀ [Austral. Mus.]. Sydney, N.S.W., 1 ♂, 1 ♀ (male forceps unarmed), [Austral. Mus.]. Minmi, N.S.W., 1 ♂ (forceps toothed). Gippsland, Victoria, XI, 10, 1919, (H. Bodley), 2 ♀.

This species is generally distributed over the tropics of the Old World and was probably originally introduced in Australia.

Labidura riparia truncata Kirby.

It is exceedingly difficult to decide whether the forms of this plastic species should be recognised as races or as mere variants. In all Australian material referable to *truncata* the ultimate tergite has (unlike *pluvialis*) a transverse caudal margin, the males do or do not have an internal tooth near the middle of the forceps and there is a tooth or swelling near their apices. This latter feature is peculiar to Australian material, so that racial recognition of *truncata* would appear to be warranted.

Mjoberg's *Labidura leucotarsata*, described from Perth, Western Australia, in 1924, is evidently a synonym.

Pawella, Queensland, (F. Bradshaw), 1 ♂ (medium, forceps with post-median tooth prominent, a very faint sub-apical swelling present), [Acad. Nat. Sci. Phila.]. Mt. Morgan District, Queensland, (Mrs. McC. Metzer) 1 ♂ (medium robust, forceps with post-median tooth prominent, a marked sub-apical swelling present), [Austral. Mus.]. Goulburn, New South Wales, 1 ♂ (slender, forceps with median tooth absent, sub-apical tooth distinct), [Austral. Mus.]. Penrith, Nepean River, N.S.W., IV, 17, 1918, (A. Musgrave), 1 ♂ (forceps with sub-median and sub-apical teeth present), [Austral. Mus.]. Port Kembla, N.S.W., III, 1927, (M. E. Higs), 1 ♂ (very robust, forceps with post-median tooth present, a very faint sub-apical swelling indicated). Waterfall, N.S.W., IV, 1928 (E. Shaw), 1 ♀, [Austral. Mus.]. Cadia near Belubula, N.S.W., V, 1923, (Miss S. E. Hosie), 1 ♀. Warrawee, Sydney, N.S.W., (H. C. Fitzhardinge), 1 ♀, [Austral. Mus.]. Arncliffe, Sydney, N.S.W., IV, 2, 1916, (B. Stoyles), 1 ♂ (slender, forceps with post-median and sub-apical teeth prominent). Lawler, Western Australia, 1 ♂ (post-median and sub-apical teeth prominent).

Only the Penrith and Lawler specimens have fully developed tegmina and wings, the latter represented by minute concealed rudiments in all the others.

APACHYIDÆ.

Apachyus queenslandicus Mjoberg.

Kuranda, Queensland, (Dodd), 1 ♂, 1 large juv. ♂, from Burr, labelled beccarii.

We are satisfied that Mjoberg was correct in describing this species. Only the inner margins of the wings are pale in this dark brown insect. The distal margin of the pygidium is slightly more produced than in *beccarii* and the forceps become gradually instead of suddenly spatulate distad.

The adult here recorded has the abdomen gradually widening distad.

Apachyus athertonensis Mjoberg.

Kuranda, Queensland, II, 4, 1925, (F. P. Dodd), 2 ♂, 4 ♀.

Mjoberg described a female, not a male as he supposed. The latter sex has the genitalia very much as in *queenslandicus*; the pygidium, however, has its distal margin armed with seven distinct points whereas in *queenslandicus* the median and lateral points have between them more numerous but very minute points.

The distinctive coloration is well described by Mjoberg. The present series agrees closely except that the entire sutural half of the wings is whitish. In a recessive example the head is reddish, not darkened caudad, and in several specimens the paler rows of latero-dorsal spots on the abdomen are fused to form broad latero-dorsal longitudinal bands.

LABIIDÆ.

Nesogaster amoenus (Stål.).

Mt. Lamington District, Northern Div., Papua, VII, 1927, (C. T. McNamara), 3 ♀ (one with short tegmina, two with long tegmina and fully developed wings).

These specimens differ from Sumatran individuals in having the tegmina immaculate and of a more metallic purplish black and the emargination following the proximo-internal projection of the forceps is more decided. Specific separation may be indicated, but males are necessary to determine this. The lateral margins of the pronotum are yellow buff, these areas broadening caudad. In all other respects close agreement with our Sumatran material is shown.

In the winged females the wing pads have a large proximo-internal spot of yellow buff.

The general appearance of this small earwig agrees closely with that of certain species of the genus *Labia*.

Nesogaster erichsoni (Dohrn).

LeGuillou's *Forficesila oceanica*, described in 1841 has been placed as a synonym of *Chelisoches morio*, with the type locality the Marquesas Islands. That author's specimen from Vavau Island, figured and named *Forficula oceanica*

by Blanchard in 1851 is a dark insect with pale wings; surely not the present species. Burr was therefore in error in suggesting *oceanica* as a synonym of *erichsoni*. We were equally at fault in 1927 in placing *pulchripes* Bormans in synonymy here.

Lithgow, New South Wales, VI, 1924, (H. E. P. Bracey), 1 ♂, [Austral. Mus.]. Melbourne, Victoria, (Edwards), 1 ♂. Launching Place, Yarra River, Vict., (Spry; under bark), 1 ♂, 1 large juv. ♀, from Burr determined as *erichsoni*.

The male pygidium agrees in its linguliform type with that of *amoenus*, but this large elongate red-brown species is otherwise very distinct and in appearance suggests certain species of the Ancistrogastrinae.

Nesogaster pulchripes (Bormans in Burr).

Kuranda, Queensland, II, 2, 1925, (F. P. Dodd), 1 ♀. McPherson Range, National Park, Queensland, XII, 18, 1926, (A. Musgrave), 1 ♂, [Austral. Mus.].

This is a distinctive species and it is unfortunate that it has been twice incorrectly placed in synonymy, under *amoenus* by Burr in 1911 and under *erichsoni* by Hebard in 1927. It agrees with the former species in size and Labiine appearance, with the latter in colour and unarmed forceps. The male pygidium, as it was originally described and also in being broader than long, is very different from the linguliform type found in those species.

Apovostox hilaris (Bormans).

Kuranda, Queensland, I, 24 and 27, 1925, (F. P. Dodd), 1 ♂, 1 ♀.

This insect agrees closely with *A. semiflavus* (Bormans) in the majority of characters used for generic separation, the more quadrate pronotum and decidedly longer second caudal tarsal joint suggesting that they may be generically distinct from the genotype, *A. pygidiatus* (Dubrony).

The very short female forceps, proximad fitting tightly about the rounded pygidium, thence with inner margins straight and attingent to the apex further suggest that *hilaris* may be found to be generically distinguishable from either of those species when the Austro-Melanesian genera are better understood.

The male agrees closely with the originally described male except that it is considerably larger, the pronotum is suffused with brown except latero-cephalad, the wings are dark brown and the feeble proximo-internal projection of the forceps is near the base instead of near the apex of the pygidium. These differences may well have specific significance, but without more material we can not determine this and the great variation shown in the present material of *Marava feae* and *Marava wallacei* indicate that extreme caution must be used in considering related Austro-Melanesian species.

Length of body ♂ 7·8, ♀ 7·9: length of forceps ♂ 2·8, ♀ 1·1 mm.

A much smaller female from Kuranda (length of body 6, length of forceps 9mm.) with pronotum dark brown, wings with a meso-proximal spot of buff and abdomen dark brown but paler towards the margins of the tergites dorsad (instead of solidly red-brown, becoming dark brown laterad and on ultimate tergite) may represent a distinct but related species.

Chætospania brunneri (Bormans).

Macpherson Range, National Park, Queensland, XII, 15, 1926, (A. Musgrave), 1 ♀, [Austral. Mus.]. Sydney, New South Wales, 1 ♂ (small, pale, stenolabic), [Austral. Mus.]. Fern Tree Gully, Victoria, IX, 1912, (Spry), 1 ♂ (eurylabic), 1 ♀, determined by Burr.

Chætospania australica (Bormans).

Kuranda, Queensland, (Dodd), 3 ♀, determined by Burr.

The coloration of this species is much more uniform and less brilliant than in *brunneri*. The female pygidium has two points between the small but sharply projecting latero-caudal angles and the forceps have a distinct flange which suddenly terminates before the moderately incurved apices.

Labia minor (Linnaeus).

Sydney, New South Wales, (C. Gibbons), 1 ♀, [Austral. Mus.].

This is undoubtedly an introduction in Australia. The species is widespread in Europe, has been introduced and become extensively established in temperate North America and has also been recorded from temperate South America and South Africa.

Marava (Burr).

Burr's *Andex* falls as a synonym of his *Marava*. Both were described in 1911, the latter having page priority.

Marava wallacei (Dohrn).

Burr noted in 1912 that *Labia grandis* Dubrony described in 1879 was a synonym of *Labia wallacei* described by Dohrn in 1864. Rehn's *Labia nigroflavida*, genotype by monotypy of *Andex* Burr, was based on a female from Cairns, Queensland, of the present species and is therefore another synonym.

In 1913 Mjoberg described *Spongiphora australica* from Kimberley, North-west Australia and in 1924 figured the insect and referred it to *Spongovostox*. Considerable series now before us convince us that that name is still another synonym, based upon a robust condition in which the male forceps are bidentate. Such material is before us from Java, normal specimens being also present in that series.

Mt. Lamington, Northern Div., Papua, IV to VII, 1927, (C. T. McNamara), 5 ♂, 10 ♀ (one female with short tegmina and no wings, other macropterous; forceps of four males unarmed, in one with a meso-distal tooth), [Austral. Mus.]. Ighibirei, New Guinea, VII to VIII, 1890, (Loria), 2 ♂ (macropterous, forceps

unarmed in one, with a meso-distal tooth in one). Stephansort, Astrolabe Bay, New Guinea, 1897, (Biro), 1 ♂, determined by Burr as *grandis* (macropterous, forceps unarmed). Siwai, Bougainville Island, X, 1, 1922, (E. O. Pockley), 1 ♂ (short tegmina, no wings, forceps with a meso-distal tooth). Ysabel Island, Solomon Islands, V, 30, 1925, (N. S. Hefferman), 1 ♀ (macropterous), [Austral. Mus.]. Kuranda, Queensland, (F. P. Dodd), 10 ♂, 10 ♀ (macropterous; male forceps unarmed in seven, with a meso-distal tooth in three).

Marava feæ (Dubrony).

This insect, though smaller and with very distinct male genitalia, agrees fully with *Marava wallacei* in all generic features. The pyriform antennal joints prevent the association with *Spongovostox* indicated by Burr in 1911.

Kuranda, Queensland, (Dodd), 1 ♂, 1 ♀, determined by Burr (tegmina short and whitish disto-laterad wings absent; male eurylabic); (F. P. Dodd), 7 ♂, 12 ♀ (macropterous, tegmina uniform black, wings buff proximo-laterad; three males eurylabic, three intermediate, one stenolabic).

CHELISOCHINÆ.

Hamaxas nigrorufus (Burr).

Burr was correct in assigning this species to *Hamaxas* in 1915, but Mjoberg repeated the original incorrect assignment to *Spongiphora* in 1924.

Mt. Lamington, Northern Div., Papua, V and VII, 1927, (C. Y. McNamara; one at light), 1 ♂, 3 ♀, [Austral. Mus.].

The limits of distribution of this species are now known to be Queensland and Hawaii.

Proreus duruoides new species. Text-figures 1 and 2.

This insect is readily separated from other Australian species by the general dark coloration, red-brown antennæ and limbs, buffy wings, abdomen with caudal margins of tergites very finely tuberculate, ultimate tergite nodulose distad and male forceps which curve outward and downward and then converging straighten out distad, these forceps very similar to those developed in males of the American genus *Doru*.

The species agrees closely with *Chelisoches*, its decidedly smaller size and general appearance indicating that it had best be referred to the very closely related *Proreus*.

Type.—♂; Kuranda, Queensland, Australia. February 3, 1925. (F. P. Dodd). [Hebard Collection Type No. 1220.]

Size medium for the genus, form slender as in the genotype, only moderately depressed. Head large, slightly broader than pronotum, moderately flattened, evenly weakly convex dorsad to very weak extensive depressions

meso-caudad of eyes on occiput, caudal margin feebly bi-convex. Eye considerably shorter than cheek. Antennæ with sixteen joints; first elongate, almost equal to combined length of second, third and fourth; third half again as long as broad, succeeding joints increasing in length, subpyriform but rapidly becoming elongate and rod-like. Pronotum glabrous, appreciably longer than wide, lateral margins very faintly divergent to the rather broadly convex caudal margin. Tegmina and wings fully developed, not keeled, glabrous. Abdomen with second gland prominent; surface microscopically very finely pitted and with caudal margins of tergites very finely tuberculate. Ultimate tergite elevated latero-caudad with two short longitudinal carinæ on each of these elevations, caudal portion conspicuously tuberculate. Pygidium vertical, rounded triangular with apex truncate, ventro-laterad before apex with a very small sharp projection on each side, much as in *Chelisoches ater* but not as pronounced. Forceps moderately stout and elongate, curving outward and downward in proximal two-fifths, thence straight convergent to incurved apices, at inner side of base thickened and terminating in a small sharp projection, ventro-internal margin from a brief distance beyond very minutely toothed to distal fifth where the shaft is horizontal, the first of these teeth more nearly a node than the others and the more distal along the edge of a lamella which gradually widens to the base of the distal portion. Penultimate sternite impresso-punctulate, these becoming coarse distad, apex broadly and shallowly emarginate. Limbs short. Caudal metatarsus pilose ventrad with two or three adjacent ventro-external spines, short but equal to combined length of the succeeding joints; second joint hirsute, very narrow, extending ventrad half the length of the third joint.

Allotype: ♀; same data as type but taken January 24, 1925. [Hebard Collection.]

Resembles the male closely, differing as follows. Ultimate tergite less elevated on each side toward caudal margin and without carinæ but more conspicuously tuberculate. Pygidium moderately convex declivent, slightly shorter, with a small blunt apical projection. Forceps moderately stout and elongate, straight then curving feebly distad to apex, simple, moderately triquetrous; internal margin moderately emarginate about pygidium, thence very feebly toothed to distal portion, no tooth more conspicuous than the others. Penultimate sternite with distal margin evenly convex.

Head, proximal portions of antennæ, abdomen and forceps shining blackish brown. Pronotum and tegmina very slightly paler, the former with narrow lateral portions which widen caudad translucent brown. Antennæ becoming gradually paler distad, chestnut in some, ferruginous in paler individuals. Wings cream colour. Limbs moderately translucent, chestnut in the darker examples, ferruginous in the others.

The measurements of three pairs of paratypes, bearing the same data as the type, and two large immature females, taken January 24 to February 4, follow those of the allotype. Length of body ♂ 10·8, ♀ 10 to 11·3; length of pronotal disk ♂ 1·63, ♀ 1·7 to 1·65; width of pronotum ♂ 1·42, ♀ 1·49 to 1·5; length of tegmen ♂ 2·48, ♀ 2·34 to 2·69; length of exposed portion of wing ♂ ·92, ♀ ·85 to ·92; length of forceps ♂ 3·8, ♀ 2·8 to 3 mm.





Text-fig. 1.—*Proreus doruoides* new species. Male. *Type*. Kuranda, Queensland, Australia. Dorsal view of ultimate tergite and forceps (greatly enlarged).

Text-fig. 2.—*Proreus doruoides* new species. Female. *Allotype*. Kuranda, Queensland, Australia. Dorsal view of ultimate tergite and forceps (same scale as Fig. 1).

CHELISOCHES Scudder.

Burr's *Kleiduchus*, described in 1911, with monotypic genotype *Forficesila australicus* Le Guillou, differs only from *Chelisoches* Scudder 1876, with genotype *Forficula morio* Fabricius, in having the tarsi more dilated. This feature is variable in the species of *Chelisoches* and insufficient to warrant the recognition of a genus. We therefore place *Kleiduchus* as a synonym of *Chelisoches*. Mjoberg questioned the validity of *Kleiduchus* in 1924.

Chelisoches australicus (Le Guillou).

Murray Island, Torres Straits, 1907, (Hedley and McCulloch), 1 ♂.

Chelisoches ater (Bormans).

Described and figured as *Psalidophora australica* by Dubrony in 1897 from Somerset, Australia, this species was named *ater* by Bormans in 1900

when it was found that it and *Forficesila australica* Le Guillou 1841 were both referable to *Chelisoches*.

Murray Island, Torres Straits, 1907, (Hedley and McCulloch), 1 ♂.

FORFICULIDÆ.

Elaunon bipartitus (Kirby).

Kuranda, Queensland, I, 29 and 30, 1925, (F. P. Dodd), 2 ♂ (one macrolabic, one delicately brachylabic), 1 ♀. Coonabarrabran, New South Wales, (H. J. Carter) 1 ♀, [Austral. Mus.]. Upper Colo River, N.S.W., X, 1922, (J. R. Kinghorn) 2 ♂, 2 ♀, [Austral. Mus.]. Uki, Tweed River, N.S.W., I, 9, 1923, (A. Musgrave), 4 ♀, [Austral. Mus.].

Forficula auricularia (Linnaeus).

Interlaken, Tasmania, I, 23 and 24, 1928, (G. P. Whitley), 2 ♂ (one moderately macrolabic, one brachylabic), [Austral. Mus.]. New Zealand, 1 ♂ (brachylabic), 1 ♀.

This common European species has been widely introduced in temperate North America and has as surely reached the Antipodes by the same means.

STENIXUS new genus.

A monotypic genus, including only *S. rhachynotus*, is here described. The very small longitudinal pronotum but large head and broad organs of flight suggest *Opisthocosmia* Dohrn, but the male abdomen with ultimate tergite narrow and both sloping and narrowing strongly caudad indicate closer affinity to *Narberia* Burr.

The remarkable tooth at each latero-cephalic angle of the pronotum readily distinguishes the genus, which is further characterized as follows.

Size medium small, form extremely slender. Head moderately tumid with sutures distinct. Antennæ with twelve segments, all very elongate and cylindrical except the minute second. Organs of flight fully developed; tegmina ample, not keeled. Abdomen with second lateral tubercle prominent and its apical glandular development distinct. Male abdomen scarcely widened meso-distad, unarmed laterad. Pygidium very small and narrow in both sexes. Forceps elongate and slender, of male nearly contiguous throughout. Limbs very slender, tarsi very small, caudal metatarsus very slightly longer than combined length of the two succeeding joints.

Stenixus rhachynotus new species. Text-figures 3 and 4.

This is not only one of the slenderest but also one of the most distinctive of the known earwigs.

Type.: ♂; Mt. Lamington, Northern Division, Papua. May, 1927. (C. T. McNamara) [Australian Museum.]

Size medium small, form extremely slender. Head with paired impressions on frons, occiput showing impression at sutures. Pronotum very small, decidedly narrower than head, considerably longer than wide; prozonal portion moderately convex to very narrow lateral portions, defined from metazona by a broadly convex sulcus; metazona narrow, feebly convex with caudal margin cingulate and broadly convex; lateral margins of pronotum cingulate, faintly converging caudad, terminating at latero-cephalic angles in a small acute-conical projection directed dorsad and weakly laterad. Tegmina decidedly broader than head, full, with caudal margin transverse. Wings very



Text-fig. 3.—*Stenixus rhachynotus* new species. Male. *Type*. Mt. Lamington, Northern Division, Papua. Dorsal view of ultimate tergite and forceps (greatly enlarged).

Text-fig. 4.—*Stenixus rhachynotus* new species. Male. *Type*. Mt. Lamington, Northern Division, Papua. Dorsal view of pronotum. (x 15)

strongly projecting. Abdomen smooth, moderately convex dorsad. Ultimate tergite simple, slightly longer than its apical width, slightly impressed meso-caudad. Pygidium minute, strongly declivent, in dorsal aspect appearing rounded and with projection equal to width. Forceps very elongate and slender, straight and subattingent to near apex with ventro-internal margin very narrowly lamellate and serrulate, this becoming gradually very weak distad, this terminating suddenly at less than distal quarter with inner surface then weakly concave to a ventro-internal tooth situated mesad in this section, thence concave due to the shaft curving inward from that point to the acute apex. Penultimate sternite broader than long, lateral margins convergent, rounding into the very feebly concave, truncate apex. Caudal tarsus only one-third as long as caudal tibia.

Allotype: ♀; same data as type. [Australian Museum.]

Very similar to male, larger. Pygidium unusually similar. Forceps proportionately shorter, attingent, straight to the unspecialized and weakly incurved apices, ventro-internal narrow lamellation similarly toothed and gradually disappearing distad, shaft mesad triquetrous. Penultimate sternite with apex convex.

Head and pronotum shining black. Tegmina and wings the same or slightly paler, deep chestnut brown. Abdomen and forceps hazel with sides darkened in one specimen. Antennæ with first two joints brown, thence buffy brown becoming gradually more and more suffused distad. Limbs clear testaceous with a very faint tawny tinge.

The measurements of a female paratype, bearing the same data, follow those of the allotype. A fragmentary paratype bearing the same data is also before us. Length of body ♂ 8·8, ♀ 10 and 10·3; length of pronotal disk (not including neck) ♂ ·92, ♀ ·92 and 1·06; greatest (cephalic) width of pronotum (not including projections the tips of which reach slightly beyond the lateral margins) ♂ ·85, ♀ ·88 and ·87; length of tegmen ♂ 2·4, ♀ 2·5 and 2·62; width across tegmina ♂ 1·77, ♀ 1·98 and 2·05; length of exposed portion of wing ♂ 1.25, ♀ 1·49, and 1·56; greatest (meso-distal) width of abdomen ♂ 1·63, ♀ 1·77 and 1·98; length of forceps ♂ 4·1, ♀ 3·8 and 3·9 mm.

Frederick Phillips, Government Printer, Brisbane.